

Research article

Open Access

## Tongue lesions in psoriasis: a controlled study

Maryam Daneshpazhoooh\*, Homayoon Moslehi, Maryam Akhyani and Marjan Etesami

Address: Department of Dermatology, Tehran University of Medical Sciences, RAZI Hospital, Vahdate-Eslami Sq. 11966 Tehran, Iran

Email: Maryam Daneshpazhoooh\* - maryamdanesh@yahoo.com; Homayoon Moslehi - moslehi\_h@yahoo.com; Maryam Akhyani - ma\_akhyani@yahoo.com; Marjan Etesami - marjanetesami@yahoo.com

\* Corresponding author

Published: 04 November 2004

Received: 17 June 2004

BMC Dermatology 2004, 4:16 doi:10.1186/1471-5945-4-16

Accepted: 04 November 2004

This article is available from: <http://www.biomedcentral.com/1471-5945/4/16>

© 2004 Daneshpazhoooh et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

### Abstract

**Background:** Our objective was to study tongue lesions and their significance in psoriatic patients.

**Methods:** The oral mucosa was examined in 200 psoriatic patients presenting to Razi Hospital in Tehran, Iran, and 200 matched controls.

**Results:** Fissured tongue (FT) and benign migratory glossitis (BMG) were the two most frequent findings. FT was seen more frequently in psoriatic patients ( $n = 66$ , 33%) than the control group ( $n = 19$ , 9.5%) [odds ratio (OR): 4.69; 95% confidence interval (CI): 2.61–8.52] ( $p$ -value  $< 0.0001$ ). BMG, too, was significantly more frequent in psoriatic patients (28 cases, 14%) than the control group (12 cases, 6%) (OR: 2.55; 95% CI: 1.20–5.50) ( $p$ -value  $< 0.012$ ). In 11 patients (5.5%), FT and BMG coexisted.

FT was more frequent in pustular psoriasis (7 cases, 53.8%) than erythematous-squamous types (56 cases, 30.4%). On the other hand, the frequency of BMG increased with the severity of psoriasis in plaque-type psoriasis assessed by psoriasis area and severity index (PASI) score.

**Conclusions:** Nonspecific tongue lesions are frequently observed in psoriasis. Further studies are recommended to substantiate the clinical significance of these seemingly nonspecific findings in suspected psoriatic cases.

### Background

The occurrence of psoriatic lesions on oral mucous membranes was a subject of controversy [1,2]. Some investigators stated that they do not occur [3]; others, have claimed that they are uncommon. Still others say that they occur only in generalized pustular psoriasis (GPP) [4,5]. Nowadays, there is sufficient evidence that a subset of patients have oral lesions in association with skin disease [2].

Oppenheim, in 1903, was the first to substantiate oral psoriasis with biopsy [6]. Since then, various lesions have

been described, including grey, yellowish, white or translucent plaques or annular forms, diffuse areas of erythema, geographic tongue and fissured tongue [7-16].

In all the cases reported in the literature, a positive biopsy showing a psoriasiform pattern has been the crucial component of the diagnosis [5,17-20]. Thus hyperkeratosis, parakeratosis, and an inflammatory infiltrate consisting of lymphocytes, polymorphonuclear leukocytes and histiocytes have been noted as well as Munro's microabscesses and spongiform pustules of Kogoj. In addition, many

**Table 1: The frequency of oral findings in psoriasis patients and control group**

	Psoriasis	Control
Fissured tongue	66	19
Benign migratory glossitis	28	12
Diffuse oral and tongue erythema	11	3
Hairy tongue	2	4
Rhomboid glossitis	2	3
Depapillated tongue	2	3

**Table 2: Frequency of fissured tongue and benign migratory glossitis according to severity in plaque-type psoriasis**

	Fissured tongue	Benign migratory glossitis
Mild	14 (26.4%)	3 (5.7%)
Moderate	23 (38.3%)	10 (16.7%)
Severe	7 (25%)	9 (32.1%)

investigators believe that the presence of cutaneous lesions with a course parallel to that of oral lesions is necessary for establishing the diagnosis of oral psoriasis [2,5,20].

However, it is impossible to perform an oral biopsy in psoriatic cases in everyday clinical practice. On the other hand, some of the lesions seen more frequently in psoriatic patients are not specific histologically. In fact, similar changes are seen in otherwise healthy people (although with a lower frequency) leading to an underestimation of the value of these findings in psoriatic patients.

In order to substantiate further the relationship between these oral disorders and psoriasis, we compared 200 patients with psoriasis to a matched control group.

**Methods**

Two hundred psoriatic patients (70 women and 130 men) attending the dermatology clinics of Razi Hospital, a major referral center in Tehran, from September 2000 till February 2001, were enrolled in this study using simple nonrandom (sequential) sampling. The diagnosis was made mainly on clinical data. The control group included 200 healthy subjects among the visitors of Surgery wards in a general hospital, matched one by one for age and sex. The skin and oral mucosa were examined in the two groups and, in addition to demographic and clinical data, PASI score [21] was recorded in plaque-type psoriasis. The data were analyzed by Epi-Info (version 6) software, and

frequency, mean, standard deviation, OR and p-value were calculated.

**Results**

The mean age of the patient group was 33.8+/- 18.2 years (4-79 years). The mean age of onset of disease was 26+/- 17.7 years (0-74 years), 23 +/-18.8 years in women and 27.6 +/- 17.0 years in men. Age and sex were matched between patients and control subjects. Family history of psoriasis was positive in 34 patients. Different clinical types of psoriasis were as follows: Chronic plaque-type psoriasis (n = 140); generalized pustular psoriasis (n = 10); flexural psoriasis (n = 10); erythrodermic psoriasis (n = 9); localized pustular psoriasis (n = 3); guttate psoriasis (n = 9); palmoplantar psoriasis (n = 15); scalp (n = 95); nail alone (n = 3).

Oral findings were detected in 87 (43.5%) and 39 (19.5%) cases in the psoriatic and control groups, respectively. They are presented in table 1. FT was seen more frequently in psoriatic patients (66 patients, 33%) than the control group (19, 9.5%) (OR: 4.69; 95% CI: 2.61-8.52) (p-value < 0.0001). BMG, too, was significantly more frequent in psoriatic patients (28 cases, 14%) than the control group (12, 6%) (OR: 2.55; 95% CI: 1.20-5.50) (p-value < 0.012). BMG was seen in 18.2% of patients with FT, and 42.9% of patients with BMG suffered from FT. In other words, in 12 patients (6%) FT and BMG coexisted. In the control group, FT and BMG coexisted in 2 cases (1%).

One hundred eighty-four patients (92%) suffered from erythematous-squamous lesions and 13 cases (6.5%) from pustular lesions. The frequency of FT in the erythematous-squamous and pustular groups was 30.4% (56 cases) and 53.8% (7 cases), respectively. On the other hand, the frequency of BMG in the erythematous-squamous and pustular groups was 14.1% (26 cases) and 15.4% (2 cases), respectively.

The severity of chronic plaque-type psoriasis cases assessed by PASI score was as follows: mild, 53 cases (37.9%); moderate, 60 cases (42.9%); and severe 27 cases (19.3%). The corresponding frequency of FT and BMG in the three severity groups is presented in table 2. The frequency of BMG increased with the severity of skin lesions (p-value < 0.001).

**Discussion**

In general oral lesions in psoriasis can be divided into two major categories. The first one includes authentic psoriatic lesions proved by biopsy and with a parallel clinical course with skin lesions. It's not known whether these lesions are truly rare, or they remain undetected, as mucosal biopsy is seldom done in known psoriatic cases.

The second group comprises the majority of oral findings in psoriasis and includes nonspecific lesions such as FT and psoriasiform lesions such as BMG [22]. These lesions are underestimated in the literature, but deserve more attention due to their high frequency.

We will discuss the main oral findings observed in our study as well as those reported in the literature.

Fissured tongue, also termed *lingua fissurata*, *lingua plicata*, *scrotal tongue*, and *grooved tongue* is recognized clinically by an antero-posteriorly oriented fissure, often with branch fissures extending laterally. It's believed by most authors to be an inherited trait. The frequency of FT increases with age and has been associated with Down's syndrome and the Melkersen-Rosenthal syndrome [5,20].

According to our study, FT was the most common oral finding in the psoriasis group: Nearly one-third of patients suffered from FT. It was significantly more frequent in psoriasis patients than the control group (9.5%) ( $p$ -value < 0.0001). The previously reported figures of the frequency of FT in the general population vary markedly in the literature depending on the study design and the target study. Axell reported a figure of 6.5% [10] and Morris found FT in 20.3% of its target study [17]. Aboyans et al reported a frequency of 2.56% in Iran [23]. On the other hand, FT was reported in 6–16.7% of psoriatic patients in different studies [3,10,14,15,20].

BMG or geographic tongue presents clinically as one or more erythematous patches with a raised white or yellow serpiginous border. Lesions may migrate across the tongue by healing on one edge while extending on another. BMG has no known cause, but it has been associated with atopic conditions, diabetes mellitus, reactive bronchitis, anemia, stress [20], hormonal disturbances, Down's syndrome and lithium therapy [24]. Lesions identical to BMG have been described in patients with Reiter's syndrome and psoriasis. The association of both psoriasis and BMG with HLA-CW6 provides further evidence that the two disorders are related [25].

In our study, BMG was significantly more frequent in psoriatic patients (14%) than the control group (6%) ( $p$ -value < 0.012). According to the literature, the estimated frequency of BMG in the general population is from 1–5% [1,10,20,26] and varies from 1–10.3% in psoriatic patients [3,10,13-15,20,26]. Only Hietanen found a figure of 1% in psoriasis [10].

FT was more frequent in patients with pustular lesions compared with the erythematous-squamous types. Contrary to previous studies, this finding was not seen for BMG, a disease generally considered accompanied with GPP

[27,28]. This may be due to the low frequency of GPP in our study group. On the other hand, the frequency of BMG increased with the severity of psoriasis in plaque-type disease, a finding not seen in Morris's study [20], perhaps due to different definition for the severity of the disease. According to our study, the frequency of FT didn't increase by increasing severity of psoriasis.

SAM was first described by Cooke in 1955 as an idiopathic inflammatory condition of the nonlingual oral mucosa [10,15]. It is also denoted using different terms: Geographic stomatitis, ectopic geographic tongue, erythema circinate migrans, and migratory stomatitis. These lesions are similar in appearance to BMG, but occur on the oral mucosal surfaces as well as the dorsum of the tongue [1,29,30].

As seen in Van der Wal's study, [14] we didn't find SAM in psoriatic patients. The reported frequency of this oral finding in psoriatic patients in the literature is between 0–19% [15,20]. Furthermore, this lesion seems to be very rare in the general population, too: Bouquot found no patients with SAM in 231616 white American dental patients [20].

Diffuse oral and tongue erythema was another positive finding in the psoriasis group with a frequency of 5.5%. This lesion, too, was reported previously in the literature, although with a lower frequency (1%) [10].

An association between FT and BMG is well established in the literature [31,32]. In our study, BMG was seen in 18.2% of patients with FT (results consistent with Pindorf's) [14,16].

## Conclusions

Overall, although oral lesions might not be considered authentic oral psoriasis unless proven histologically and with a parallel clinical course, nonspecific tongue lesions are significantly more frequent in psoriatic cases. Further studies are recommended to evaluate the clinical significance of these seemingly nonspecific lesions in a suspected psoriatic case. Furthermore, more thorough studies are recommended regarding the relationship of oral psoriasis and disease severity in plaque-type psoriasis.

## Competing interests

The author(s) declare that they have no competing interests.

## Authors' contributions

All authors contributed equally in the study design, literature search, data analysis and manuscript preparation. All authors read and approved the final manuscript.

## References

- Weathers DR, Baker G, Archard HO, Burkes EJ: **Psoriasiform lesions of the oral mucosa (with emphasis on "ectopic geographic tongue")**. *Oral surgery* 1974, **37**:812-888.
- Bruce AJ, Roger RS III: **Oral psoriasis**. *Dermatol Clin* 2003, **21**:99-104.
- Buchner A, Begleiter A: **Oral lesions in psoriatic patients**. *Oral Surg Oral Med Oral Pathol* 1976, **41**:38-40.
- Trigonides G, Markopoulos AK, Konstantinidis AB: **Dermal psoriasis involving the oral cavity**. *J Oral Med* 1986, **41**:98-101.
- Younai SF, Phelan JA: **Oral mucositis with features of psoriasis: Report of a case and review of the literature**. *Oral Surg Oral Med Oral Pathol* 1997, **84**:61-7.
- Fischman SL, Barnett ML, Nisengard RJ: **Histopathologic, ultrastructural, immunologic findings in an oral psoriatic lesion**. *Oral Surg Oral Med Oral Pathol* 1977, **44**:253-60.
- Brice DM, Danesh-Meyer MJ: **Oral lesions in patients with psoriasis: Clinical presentation and management**. *J Periodontol* 2000, **71**:1896-1903.
- Richardson LJ, Kratochvil FJ, Zieper MB: **Unusual palatal presentation of oral psoriasis**. *J Can Dent Assoc* 2000, **66**:80-2.
- Scully C: **The oral cavity**. In *In Rook/ Wilkinson/ Ebling Textbook of dermatology* 6th edition. Edited by: Champion RH, Burton JL, Burns DA, Breathnach SM. Oxford: Blackwell Science Ltd; 1998:3047-3123.
- Hietanen J, Salo OP, Kanerva L, Juvakoski T: **Study of the oral mucosa in 200 consecutive patients with psoriasis**. *Scand J Dent Res* 1984, **92**:50-4.
- O'Keefe E, Braverman IM, Cohen I: **Annulus migrans. Identical lesions in pustular psoriasis, Reiter's syndrome, and geographic tongue**. *Arch Dermatol* 1973, **107**:240-4.
- Odom RB, James WD, Berger TG: *Andrew's diseases of the skin. Clinical dermatology* Ninth edition. WB. Saunders Company. Philadelphia; 2000:998.
- Ulmansky M, Michelle R, Azaz B: **Oral psoriasis: report of six new cases**. *J Oral Pathol Med* 1995, **24**:42-5.
- Van der Wal N, van der Kwast WAM, van Dijk E, van der Waal I: **Geographic stomatitis and psoriasis**. *Int J Oral Maxillofac Surg* 1998, **17**:106-9.
- Pogrel MA, Cram D: **Intraoral findings in patients with psoriasis with special reference to ectopic geographic tongue (erythema circinata)**. *Oral Surg Oral Med Oral Pathol* 1976, **41**:174-81.
- Dawson TAJ: **Tongue lesions in generalized pustular psoriasis**. *Br J Dermatol* 1974, **91**:419-24.
- Salmon TA, Robertson GR, Tracy NH, Hiatt WR: **Oral psoriasis**. *Oral Surg Oral Med Oral Pathol* 1974, **38**:48-54.
- Jones LE, Dolby AE: **Desquamative gingivitis associated with psoriasis**. *J Periodontol* 1972, **43**:35-7.
- DeGregori G, Pippen R, Davies E: **Psoriasis of the gingival and the tongue: Report of a case**. *J Periodontol* 1971, **42**:97-100.
- Morris LF, Phillips CM, Binnie WH, Sander HM, Silverman AK, Menter MA: **Oral lesions in patients with psoriasis: a controlled study**. *Cutis* 1992, **49**:339-344.
- El Sayed MF, Marghuery M: **Psoriasis**. *Ann Dermatol Venereol* 1997, **124**:91-104.
- Femiano F: **Geographic tongue (migrant glossitis) and psoriasis**. *Minerva Stomatol* 2001, **50**:213-7.
- Aboyans V, Ghaemmaghami A: **The incidence of fissured tongue among 4009 Iranian dental outpatients**. *Oral Surg Oral Med Oral Pathol* 1973, **36**:34-8.
- Assimakopoulos D, Patikakos G, Fotika C, Elisaf M: **Benign migratory glossitis or geographic tongue: an enigmatic oral lesion**. *Am J Med* 2002, **113**:751-755.
- Camp RDR: **Psoriasis**. In *In Rook/ Wilkinson/ Ebling Textbook of dermatology* 6th edition. Edited by: Champion RH, Burton JL, Burns DA, Breathnach SM. Oxford. Blackwell Science Ltd; 1998:1589-1649.
- Robinson CM, Di Biase AT, Leigh IM, Williams DM, Thornhill M: **Oral psoriasis**. *Br J Dermatol* 1996, **134**:347-9.
- Wagner G, Luckasen JR, Goltz RW: **Mucous membrane involvement in generalized pustular psoriasis. Report of three cases and review of the literature**. *Arch Dermatol* 1976, **112**:1010-14.
- Baker H, Ryan TJ: **Generalized pustular psoriasis. A clinical and epidemiologic study of 104 cases**. *Br J Dermatol* 1968, **80**:771-93.
- Zunt SL, Tomich CE: **Erythema migrans - a psoriasiform lesion of the oral mucosa**. *J Dermatol Surg Oncol* 1989, **15**:1067-1070.
- Ralls SA, Warnock GR: **Stomatitis areata migrans affecting the oral cavity**. *Oral Surg Oral Med Oral Pathol* 1985, **60**:197-200.
- White DK, Leis HJ, Miller AS: **Intraoral psoriasis associated with widespread dermal psoriasis**. *Oral Surg Oral Med Oral Pathol* 1976, **4**:174-181.
- Eidelman E, Chosack A, Cohen T: **Scrotal tongue and geographic tongue: polygenic and associated traits**. *Oral Surg Oral Med Oral Pathol* 1976, **42**:591-596.

## Pre-publication history

The pre-publication history for this paper can be accessed here:

<http://www.biomedcentral.com/1471-5945/4/16/prepub>

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
[http://www.biomedcentral.com/info/publishing\\_adv.asp](http://www.biomedcentral.com/info/publishing_adv.asp)

