

Research article

Joint effects of citrus peel use and black tea intake on the risk of squamous cell carcinoma of the skin

Iman A Hakim* and Robin B Harris

Address: University of Arizona College of Public Health and the Arizona Cancer Center, 1515 N Campbell Ave, Tucson, AZ 85724, USA

E-mail: Iman A Hakim* - ihakim@azcc.arizona.edu; Robin B Harris - rharris@azcc.arizona.edu

*Corresponding author

Published: 1 August 2001

Received: 9 April 2001

BMC Dermatology 2001, 1:3

Accepted: 1 August 2001

This article is available from: <http://www.biomedcentral.com/1471-5945/1/3>

© 2001 Hakim and Harris; licensee BioMed Central Ltd. Verbatim copying and redistribution of this article are permitted in any medium for any non-commercial purpose, provided this notice is preserved along with the article's original URL. For commercial use, contact info@biomedcentral.com

Abstract

Background: Differences in tea drinking habits and/or citrus peel use are likely to vary by populations and could contribute to the inconsistencies found between studies comparing their consumption and cancer risk.

Methods: A population-based case-control study was used to evaluate the relationships between citrus peel use and black tea intake and squamous cell carcinoma (SCC) of the skin. Moreover, we assessed the independent and interactive effects of citrus peel and black tea in the development of SCC.

Results: Hot and iced teas were consumed by 30.7% and 51.8% of the subjects, respectively. Peel consumption was reported by 34.5% of subjects. Controls were more likely than were cases to report citrus peel use (odds ratio (OR) = 0.67) and hot tea intake (OR = 0.79). After adjustment for hot and iced tea intake, the ORs associated with citrus peel use were 0.55 and 0.69, respectively, whereas the corresponding adjusted ORs for hot and iced tea intake after adjustment for citrus peel use were 0.87 and 1.22 respectively. Compared with those who did not consume hot black tea or citrus peel, the adjusted ORs associated with sole consumption of hot black tea or citrus peel were 0.60 and 0.30, respectively. Subjects who reported consumption of both hot black tea and citrus peel had a significant marked decrease (OR= 0.22; 95% CI = 0.10 – 0.51) risk of skin SCC.

Conclusion: These results indicate that both citrus peel use and strong (hot) black tea have independent potential protective effects in relation to skin SCC.

Introduction

Non-melanoma skin cancer, which includes squamous cell carcinoma (SCC) and basal cell carcinoma (BCC), is the most common type of cancer in the United States and results in substantial morbidity and treatment costs. It is estimated that 1.2 million new cases of non-melanoma skin cancer occur each year in the United States, and the majority of these are believed to result from heavy expo-

sure to ultraviolet (UV) light from the sun [1]. The incidence of skin cancer is predicted to increase even further because of the depletion of the stratospheric ozone layer [2]. The development of chemopreventive strategies for skin cancer is thus a high public health priority. Residents of Arizona experience a three to seven times greater incidence of non-melanoma skin cancer than the general population in the United States [3]. Our goal is

the development of oral agents that will complement primary skin cancer prevention.

Black tea has been shown to have a preventative effect against skin carcinogenesis in animal models [4–9]. Of particular interest are the accumulating data that reflect similar potential health benefits associated with both the epicatechins of green tea and the theaflavins of black tea [10,11]. Several mechanisms may be responsible for the anti-initiation and anti-promotion properties of black tea [5,10–15]

Citrus, in addition to providing an ample supply of vitamin C, folic acid, potassium, and pectin, contains a host of active phytochemicals. The mutagenicity-reducing activity per weight of peels of citrus fruits was considerably higher than that of their juices [16]. The two main compositional differences between peel and juice components are that the peel contains a higher concentration of ascorbic acid than the juice, and that the peel also contains higher concentrations of active components (d-limonene, hesperidin, naringin, and auraptene) than do the juice and pulp.

d-Limonene, which comprises > 90% of citrus peel oil, has demonstrated chemopreventive activity against a variety of chemically induced rodent cancers [17–22]. Several mechanisms of action may account for the antitumor activities of d-limonene [23–25]. Among citrus bioflavonoids, hesperidin and naringin have been tested as potential chemopreventive agents [26,27]. In our previous studies, we showed that citrus peel use [28] and consumption of hot (strong) black tea [29] were each independently associated with reduced risk of skin SCC. Neither study, however, explored the potential for a joint or synergistic effect of tea and citrus consumption. Therefore, the present study sought to examine the independent and interactive roles of citrus peel use and black tea consumption in skin SCC in Arizona, a population at high risk of skin SCC.

Subjects & Methods

Study population

Subjects and methods of the Southeastern Arizona Health Study (SEAHS) have been described previously [28,29]. Briefly, SEAHS is a population-based case-control study that was conducted from 1994–1996 with cases of SCC of the skin randomly selected from persons identified through the Southeastern Arizona Skin Cancer Registry. Cases were eligible if they were ≥ 30 years of age, had a histo-pathologically confirmed, non-metastatic SCC of the skin diagnosed within the past 4 months, with no prior history of a skin cancer. Control subjects were recruited through a random-digit dialing process to reflect the age and gender distribution of the cases. One

control per household was invited to participate using modified Waksberg criteria [30]. Control subjects were considered eligible if they had no prior history of skin cancer, lived within the Tucson region, and were within the age, gender, and ethnicity grouping. All subjects (404 cases and 391 controls) completed an extensive interview for demographic, behavioral, and past ultraviolet exposure information. A subset of the total study population ($n = 566$), namely those who also completed 24-hour dietary recalls, were asked to complete the citrus and tea consumption questionnaires, yielding 450 subjects (234 cases and 216 controls) in this study (tea & citrus).

Citrus & tea questionnaires

Citrus (CQ) and tea (TQ) questionnaires were developed after a series of focus groups. Both questionnaires were tested for short and long term reliability as well as relative validity (diet methods). CQ asked detailed information on consumption of each type of citrus fruit and juice. In addition, the questionnaire asked how the juice was stored, i.e. types of containers, and whether citrus juices and citrus peel were added during food preparation and/or food serving [28]. TQ asked about usual tea intake over the past year, as well as a lifetime consumption pattern and how the past year intake differed from the lifetime pattern (amount or type or preparation techniques). Detailed information was collected for the past year's tea intake for each type of tea consumed (black, green or herbal and hot or iced) [29].

Other variables

All participants completed a structured interview detailing personal, behavioral, and demographic characteristics. This interview instrument sought information on: skin characteristics, sunburns and tanning history, use of suntan lotions and sunscreens, residential history, UV exposure during past year, family history of skin cancer, past medical history, tobacco and alcohol use, physical characteristics, and demographic information. All subjects also completed four 24-hour dietary recalls, which included 4 randomly selected days within two weeks of the clinic visit. Daily mean nutrient intakes were calculated with the use of the Minnesota Nutrition Data System [31].

Quality control

All interviews were conducted by trained, experienced interviewers. One interviewer conducted over 90% of the personal interviews and 100% of the dietary recalls, and citrus and tea questionnaires. At the time of CQ and TQ, the interviewer was unaware of the case-control status of the subjects. She was given a computer-generated list that included only names and phone numbers of subjects. After each interview, questionnaires were reviewed for completeness and coded. Data entry was through

screen-based entry programs that included range checks.

Data analyses

Distribution of demographic characteristics and potential risk factors were compared between cases and controls using *t* tests for continuous variables and Chi-square (X^2) tests for categorical variables. Chi-square (X^2) tests for trend were also calculated. To assess the risk of skin SCC with citrus peel use and black tea intake, crude and adjusted odds ratios (ORs) with their 95 percent confidence intervals (C.I.) were estimated with logistic regression models.

We assessed the potential confounding effects of age, sex, education, energy intake (kcal/day), fat intake (mean percent of kcal as fat), retinol and carotenoids ($\mu\text{g}/\text{day}$), vitamin C (mg/day), α -tocopherol (mg/day), foods rich in carotenoids (servings/week), alcohol intake (mean alcohol intake/day), smoking history (never, former smoker, and current smoker), body mass index (kg/ht in m^2), daily hours of sun exposure during the past year, history of actinic skin damage (self-reported physician-diagnosed actinic keratosis), self-reported ability to tan after prolonged sun exposure (no suntan, mildly tan, moderately tan, and deeply tan), and number of current freckles on the arms. The initial adjusted model included age, sex. Inclusion of energy intake, fat intake, antioxidant vitamins (vitamin A, E and C) or carotenoid intake, alcohol intake, smoking status, education, and daily hours of sun exposure did not alter any of the results and were excluded from the final model.

Age, sex, inability to tan after prolonged sun exposure, number of current freckles on the arms, and history of diagnosed and treated actinic keratosis (AK) were included in the final multivariate model. All statistical analyses were done using STATA computer software [32].

Results

A total of 263 men and 187 women who had participated in the SEAHs are included in this current study. Since this evaluation of citrus peel and tea consumption in relation to skin SCC was completed in a subset of the original participants, we compared subjects who participated in this study with those who did not participate. We found no statistically significant difference between the two groups in relation to case-control status, gender, education, smoking, average hours in sun per day in past year, and tanning ability. Participants in the current study did report more AK history than subjects who did not participate in the study, with the increase being consistent for cases and controls.

Table 1 shows the distribution of cases and controls according to sex, age, smoking status, alcohol intake, reported tanning ability, history of AK, number of current freckles on the arms, and daily hours of sun exposure during the past year. Information from CQ and TQ, as citrus peel use, hot and iced black tea consumption, are also included. The study population is an older educated Southwestern United States population with 68.8% of cases and 66.2% of controls having some college education. There was no difference between the cases and controls in antioxidants intake (retinol in $\mu\text{g}/\text{day}$, β -carotene in $\mu\text{g}/\text{day}$, α -carotene in $\mu\text{g}/\text{day}$, lycopene in $\mu\text{g}/\text{day}$, α -tocopherol in mg/day, and vitamin C in mg/day). There were also no differences between the cases and controls in the number of years they have lived in Arizona, and in the reported usual hours spent in the sun. Cases reported spending in the past year, on average, 1.38 hours in the sun during peak hours compared to 1.49 hours for controls. Only tanning ability, history of AK, number of current freckles on the arms, citrus peel use and hot black tea consumption showed a significant difference between cases and controls.

In this Arizona population, 66.4% and 34.5% reported some tea drinking and some citrus peel use, respectively, during the past year. Controls were more likely than were cases to report citrus peel use (39.2% versus 30.2%, with an OR of 0.67) and hot tea intake (35.7% versus 26.1%, with an OR of 0.79).

Table 2 shows the ORs for citrus peel use and tea intake controlling for the other variables. The OR of skin SCC associated with any citrus peel use was 0.67 (95%CI = 0.45 – 0.99). Adjustment for hot tea intake resulted in a decrease in the OR to 0.55 (95%CI = 0.34 – 0.91), while adjustment for iced tea consumption did no change the OR (OR = 0.69; 95%CI = 0.45 – 1.05). Further adjustment for age, gender, AK, tanning ability and number of current freckles on the arms resulted in marked decrease in ORs to 0.34 (95%CI = 0.18 – 0.62) and 0.54 (95%CI = 0.33 – 0.89) among hot and iced tea drinkers respectively.

Taking non-tea drinkers as the reference group, the OR of skin SCC was 0.79 (95% CI = 0.49 – 1.26) for drinkers of hot tea compared to 1.14 (95%CI = 0.76 – 1.72) for drinkers of iced tea. After adjustment for citrus peel use, the ORs increased to 0.87 (95%CI = 0.54 – 1.39) and 1.22 (95% CI = 0.81 – 1.84) for drinkers of hot and iced tea respectively. However, in the final model the ORs of skin SCC decreased to 0.65 (95%CI = 0.37 – 1.15) and 1.06 (95%CI = 0.37 – 1.15) among hot and iced tea drinkers respectively.

Table 1: Selected characteristics of skin SCC cases and control participating in SEAH study

	Cases N = 234 Mean + SD or %	Controls N= 216 Mean + SD or %	p
Age (mean + SD)	66.6 + 10.4	66.2 + 11.1	0.75
Male (%)	59.0%	57.9%	0.81
Alcohol ^a	Yes 78.7%	81.8%	0.53
Smoking :	Never 36.8%	37.5%	
	Former 51.7%	46.8%	
	Current 11.5%	15.7%	0.36
Hours in Sun ^d (h/day)	1.38 + 0.1	1.49 + 0.1	0.35
AK ^a	Yes 78.2%	37.0%	0.000
Tanning ^b	Yes 68.8%	81.0%	0.001
Freckles ^c	No 50.6%	61.5%	0.005
Citrus peel use:	Yes 30.2%	39.2%	0.045
Hot tea intake:	Yes 26.1%	35.7%	0.028
Iced tea intake	Yes 53.4%	50.0%	0.47
Kcal/day (mean + SD)	1499.9 + 396.3	1531.8 + 433.5	0.41
Antioxidants (mean + SD)			
Retinol (µg/day)	474.5 + 23.2	449.4 + 23.8	0.45
β-Carotene (µg/day)	92.4 + 7.1	87.6 + 7.8	0.65
α-Carotene (µg/day)	722.9 + 52.3	777.9 + 63.0	0.50
Lycopene (µg/day)	5882.9 + 359.7	5440.2 + 419.4	0.42
α-Tocopherol (mg/day)	7.1 + 1.1	6.4 + 0.5	0.57
Vitamin C (mg/day)	110.7 + 4.5	109.5 + 4.0	0.84

a: Based on 4–24 hour dietary recalls; b : History of treated actinic keratosis c: Moderate to deep tanning ability after prolonged sun exposure; d: Number of current freckles on the arms; e: Average hours per day during past year spent outside during peak sun expressed as mean (SD).

Table 2: Odds ratios and 95% confidence intervals from multivariate analyses for skin SCC

Model	Citrus peel OR (95% C.I.)	Black tea OR (95% C.I.)
Hot Black tea		
- Citrus peel	0.67 (0.45 – 0.99)	—
- Hot tea	—	0.79 (0.49 – 1.26)
- Citrus peel + hot tea	0.55 (0.34 – 0.91)	0.87 (0.54 – 1.39)
- The above + age & gender	0.54 (0.32 – 0.90)	0.84 (0.52 – 1.36)
- The above + SCC risk factors*	0.34 (0.18 – 0.62)	0.65 (0.37 – 1.15)
Iced Black tea		
- Citrus peel	0.67 (0.45 – 0.99)	—
- Iced tea	—	1.14 (0.76 – 1.72)
- Citrus peel + iced tea	0.69 (0.45 – 1.05)	1.22 (0.81 – 1.84)
- The above + age & gender	0.69 (0.45 – 1.07)	1.21 (0.80 – 1.84)
- The above + SCC risk factors*	0.54 (0.33 – 0.89)	1.06 (0.37 – 1.15)

* History of treated actinic keratosis, tanning ability after prolonged sun exposure and number of current freckles on the arms.

The independent and combined effects of citrus peel and hot black tea consumption are shown in Table 3. Among those who did not consume citrus peel, 38.8% (38/98) of

the cases compared to 44.3% (39/88) of controls consumed hot black tea (p = 0.44). Whereas among those who did report citrus peel use, 59.0% (23/39) of cases

Table 3: Independent and interactive effects of hot black tea consumption and citrus peel use on the risk of skin SCC

Intake Citrus peel factors	Hot black tea	Cases No.	Controls No	Crude	OR (95% confidence Interval)	
					Adjusted for age & gender	Adjusted for age, gender & SCC risk
no	no	60	49	1	1	1
no	yes	38	39	0.80 (0.44 – 1.42)	0.75 (0.41 – 1.37)	0.60 (0.30 – 1.23)
yes	no	16	27	0.48 (0.23 – 0.99)	0.45 (0.22 – 0.97)	0.30 (0.13 – 0.72)
yes	yes	23	38	0.49 (0.26 – 0.94)	0.47 (0.24 – 0.92)	0.22 (0.10 – 0.51)
P for trend				0.013	0.012	0.000

compared to 58.5% (38/65) of controls consumed hot black tea (p = 0.96). On average, 21% of the subjects reported consumption of both citrus peel and hot tea (16.8% in cases vs 24.8% controls).

Compared with those who did not consume hot black tea or citrus peel, the adjusted ORs associated with sole consumption of hot black tea or citrus peel were 0.60 (95% CI = 0.30 – 1.23) and 0.30 (95% CI = 0.13 – 0.72), respectively. Subjects who reported consumption of both hot black tea and citrus peel had a significant marked decrease in risk of skin SCC. (OR= 0.22; 95% CI = 0.10 – 0.51).

Similarly, the independent and combined effects of citrus peel and iced black tea consumption are shown in Ta-

ble 4. Among those who did not consume citrus peel, 56.5% (78/138) of the cases compared to 55.9% (62/111) of controls consumed iced black tea (p = 0.92). Whereas among those who did report citrus peel use, 74.6% (47/63) of cases compared to 62.5% (45/72) of controls consumed iced black tea (p = 0.13). On average, 24% of the subjects reported consumption of both citrus peel and iced tea (20.4% in cases vs 24.6% controls).

Compared with those who did not consume iced black tea or citrus peel, the adjusted ORs associated with sole consumption of iced black tea and citrus peel were 0.91 (95% CI = 0.46 – 1.45) and 0.32 (95% CI = 0.14 – 0.75), respectively. Subjects who reported consumption of both iced black tea and citrus peel had lower risk of skin SCC (OR= 0.58; 95% CI = 0.30 – 1.12).

Table 4: Independent and interactive effects of iced black tea consumption and citrus peel use on the risk of skin SCC

Intake Citrus Peel	Iced black tea	Cases No.	Controls No.	Crude	OR (95% confidence Interval)	
					Adjusted for age & gender	Adjusted for age, gender & SCC risk factors
no	no	60	49	1	1	1
no	yes	78	62	1.02 (0.62 – 1.69)	1.00 (0.60 – 1.67)	0.91 (0.46 – 1.45)
yes	no	16	27	0.48 (0.23 – 0.99)	0.47 (0.22 – 0.99)	0.32 (0.14 – 0.75)
yes	yes	47	45	0.85 (0.49 – 1.49)	0.84 (0.48 – 1.50)	0.58 (0.30 – 1.12)
P for trend				0.29	0.30	0.04

Discussion

Arizona has one of the highest risks of skin SCC worldwide and tea beverages, as well as, citrus products are commonly consumed by Arizonans. This older Arizona population offered a unique opportunity to study poten-

tial associations between consumption of tea and/or citrus peel and risk of skin SCC. In this population, two-thirds of all subjects reported black tea consumption in the previous year compared to about one third reporting citrus peel use. Our data showed that persons without

skin cancer significantly consumed more citrus peel and hot black tea than did cases of skin SCC. We have previously demonstrated that among Arizonans [29], black tea beverages were prepared and consumed in various ways and that iced tea beverages were more likely to be prepared diluted.

Laboratory investigations support the capability of citrus peel or its major constituents, d-limonene, and also the ability of tea extract or its constituents, tea polyphenols to reduce not only the incidence and tumor size of chemically-induced tumors at several sites [4,8,9,17,18,20,22] but also the growth of various tumor cells in culture [11,33]. Both d-limonene, representative of citrus peel, and tea polyphenols, representative of black tea, are known to impact antiproliferative effects, including apoptosis, on neoplastic cells [4,34–37]. Dietary d-limonene has been shown to be effective in the chemoprevention and chemotherapy of cancer. As a result, its cancer chemotherapeutic activities are under evaluation in Phase I human clinical trials [38,39]. Moreover, much research has been focused on the potential use of flavonoids (in citrus and tea) as inhibitors of neoplastic transformation [40] and as free radical scavengers to prevent oxidative skin damage [41,42]

Oxidative damage is one of the many mechanisms leading to skin cancer and other chronic diseases. Black tea is a good source of antioxidants and addition of lemon to tea increases its antioxidant potential [43]. Our data indicates that among older Arizonans, citrus peel use and hot black tea intake were each independently associated with reduced risk of skin SCC. However, among these study subjects, citrus peel use was associated with a more than 70% reduced risk for skin SCC, whereas hot black tea consumption alone conferred 40% reduced risk for skin SCC. Moreover, the present study suggested that a dietary pattern that includes the consumption of both hot black tea and citrus peel markedly decreased the risk of skin SCC (ORs = 0.22 – 0.58). Even though iced black tea consumption alone was not associated with reduced risk of skin SCC (OR = 0.91), consumption of both iced black tea and citrus peel was associated with lower risk of skin SCC (OR = 0.58). Among Arizonans, citrus peel was reported to be frequently consumed along with iced tea beverages at meals.

Some limitations and strengths of the study deserve consideration. In case-control studies, the possibilities for recall and interviewer bias are a major concern. Differential recall of tea and citrus consumption between cases and controls can lead to biased estimates of effect. Furthermore, since there was a lag between diagnosis of the skin cancer and interview (average of 4 months), there is potential for cases to have altered their behavior and to

then report their recently changed behaviors. Several steps were taken to reduce potential bias. Standard questionnaires were administered to all subjects by a trained interviewer who was not aware of the case-control status of the subjects at the time of the tea and citrus questionnaire administration. There is some evidence, however, that skin cancer cases did recently alter their behavior for risk factors they thought were related to skin cancer. For instance, they reported similar recent (past year) exposure to the sun to the controls. The interview did not record information on sun exposure experiences in the more distant past; however, given that skin SCC is related to other measures of high UV exposure (i.e. history of actinic keratosis, history of sunburns), then the lack of a finding for a differential sun exposure history for cases and controls argues for a change in behavior since the diagnosis of the skin cancer.

However, while it appears that the cases did modify some of their behaviors, there is no evidence that they recently altered their consumption of tea or citrus products. Also, since there have been no prior studies of skin cancer occurrence and tea or citrus consumption, it is unlikely that this population would have considered tea or citrus peel consumption, or lack of consumption, to be related to their risk of skin cancer. Public perception of tea and citrus consumption has been that there would be no difference in the potential effect of tea or citrus based on different products.

This study controlled for important confounding factors, including history of physician-diagnosed actinic keratosis, tanning ability, and number of current freckles on the arms. Consumption of citrus peel along with black tea appeared to be generally protective. Significant independent inverse associations between citrus peel use and skin SCC were observed. Moreover, a synergistic and/or additive effect(s) on skin SCC risk was observed when both citrus peel and black tea were consumed. While this study was only of moderate sample size, particularly when exploring relationships within subgroups of different consumption patterns, we were able to observe a significant protective effect for citrus peel and hot black tea consumption in relation to skin SCC.

Competing interests

Have you in the past five years received reimbursements, fees, funding, or salary from an organization that may in any way gain or lose financially from the publication of this paper?

As cited in the acknowledgement section above: The development of the tea questionnaire was supported by a grant from Unilever Health Institute, Vlaardingen,

UNC, PO Box 114, 3130 AC Vlaardingen, The Netherlands

Do you hold any stocks or shares in an organization that may in any way gain or lose financially from the publication of this paper? No

Do you have any other financial competing interests? No

Are there any non-financial competing interests you would like to declare in relation to this paper? No

Acknowledgements

This research was supported by Public Health Service Grant K07 CA76008. Also, this publication was made possible by Public Health Service Grant P01 CA27502. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the National Cancer Institute. The development of the tea questionnaire was supported by a grant from Unilever Health Institute, Vlaardingen, UNC, PO Box 114, 3130 AC Vlaardingen, The Netherlands. The authors would like to thank Dr. David Alberts, director of the Division of Cancer Prevention and Control for his support, Steve Rodney for his assistance with data management, and Mary Lurie for her assistance with interviewing and data entry.

References

- Miller DL, Weinstock MA: **Nonmelanoma skin cancer in the United States: Incidence.** *J. Am. Acad. Dermatol.* 1994, **30**:774-778
- Scotto J, Fears TR, Fraumeni JF Jr: **Solar radiation.** In *Cancer Epidemiology and Prevention*, 2nd ed, D Schottenfeld and JF Fraumeni, Jr (eds). New York: Oxford University Press. 1996:355-372
- Robinson JK: **Risk of developing another basal cell carcinoma.** *Cancer.* 1987, **60**:118-120
- Wang ZY, Huang MT, Lou YR, Xie JG, Reuhl KR, Newmark HL, Ho CT, Yang CS, Conney AH: **Inhibitory effects of black tea, green tea, decaffeinated black tea, and decaffeinated green tea on ultraviolet B light-induced skin carcinogenesis in 7,12-dimethylbenz [a] anthracene-initiated SKH-I mice.** *Cancer Research.* 1994, **54**(13):3428-3435
- Lu YP, Lou YR, Xie JG, Yen P, Huang MT, Conney AH: **Inhibitory effect of black tea on the growth of established skin tumors in mice: effects on tumor size, apoptosis, mitosis and bromodeoxyuridine incorporation into DNA.** *Carcinogenesis.* 1997, **18**(11):2163-2169
- Lo YR, Lu YP, Xie JG, Huang MT, Conney AH: **Effects of oral administration of tea, decaffeinated tea, and caffeine on the formation and growth of tumors in high-risk SKH-I mice previously treated with ultraviolet B light.** *Nutrition & Cancer.* 1999, **33**(2):146-153
- Record IR, Dreosti IE: **CSIRO Division of Human Nutrition, Adelaide, South Australia, Australia. Protection by black tea and green tea against UVB and UVA + B induced skin cancer in hairless mice.** *Mutation Research.* 1998, **422**(1):191-199
- Katiyar SK, Mukhtar H: **Inhibition of phorbol ester tumor promoter 12-tetra-decanoylphorbol-13-acetate-caused inflammatory responses in SENCAR mouse skin by black tea polyphenols.** *Carcinogenesis.* 1997, **18**(10):1911-1916
- Javed S, Mehrotra NK, Shukla Y: **Chemopreventive effects of black tea polyphenols in mouse skin model of carcinogenesis.** *Biomedical & Environmental Sciences.* 1998, **11**(4):307-313
- Dong Z, Ma W, Huang C, Yang CS: **Inhibition of tumor promoter-induced activator protein 1 activation and cell transformation by tea polyphenols, (-)-epigallocatechin gallate, and theaflavins.** *Cancer Research.* 1997, **57**(19):4414-4419
- Liang YC, Chen YC, Lin YL, Shiau SY, Ho CT, Lin JK: **Suppression of extracellular signals and cell proliferation by the black tea polyphenol, theaflavin-3,3'-digallate.** *Carcinogenesis.* 1999, **20**(4):733-736
- Halder J, Bhaduri AN: **Protective role of black tea against oxidative damage of human red blood cells.** *Biochemical & biophysical Research Communications.* 1998, **244**(3):903-907
- Wei H, Zhang X, Zhao JF, Wang ZY, Bickers D, Leibold M: **Scavenging of hydrogen peroxide and inhibition of ultraviolet light-induced oxidative DNA damage by aqueous extracts from green and black teas.** *Free Radical Biology & Medicine.* 1999, **26**(11-12):1427-1435
- Steele VE, Bagheri D, Balentine DA, Boone CW, Mehta R, Morse MA, Sharma S, et al: **Preclinical efficacy studies of green and black tea extracts.** *Proceedings of the Society for Experimental Biology and Medicine.* 1999, **220**(4):210-212
- Chung JY, Huang C, Meng X, Dong Z, Yang CS: **Inhibition of activator protein 1 activity and cell growth by purified green tea and black tea polyphenols in H-ras-transformed cells: Structure-activity relationship and mechanisms involved.** *Cancer Research.* 1999, **59**(18):4610-4617
- Higashimoto M, Yamato H, Kinouchi T, Ohnishi Y: **Inhibitory effects of citrus fruits on the mutagenicity of 1-methyl-1,2,3,4-tetrahydro-carboline-3-carboxylic acid treated with nitrite in the presence of ethanol.** *Mutation Research/Genetic Toxicology and Environmental Mutagenesis.* 1998, **415**:219-226
- Elson CE, Maltzman TH, Boston JL, et al: **Anti-carcinogenic activity of d-limonene during the initiation and promotion/progression stages of DMBA-induced rat mammary carcinogenesis.** *Carcinogenesis.* 1988, **9**:331-332
- Maltzman TH, Hurt LM, Elson CE, Tanner MA, Gould MN: **The prevention of nitrosomethylurea-induced mammary tumors by d-limonene and orange oil.** *Carcinogenesis.* 1989, **10**:781-783
- Giri RK, Parija T, Das BR: **D-Limonene chemoprevention of hepatocarcinogenesis in AKR mice: inhibition of c-jun and c-myc.** *Oncology Reports.* 1999, **6**(5):1123-1127
- Wattenberg LW, Coccia JB: **Inhibition of 4-(methylnitrosoamino)-1-(3-pyridyl)-1-butanone carcinogenesis in mice by d-limonene and citrus fruit oils.** *Carcinogenesis.* 1991, **12**:115-117
- Yano H, Tatsuta M, Iishi H, Baba M, Sakai N, Uedo N: **Attenuation of d-limonene of sodium chloride-enhanced gastric carcinogenesis induced by N-methyl-N'-nitro-N-nitrosoguanidine in Wistar rats.** *International journal of cancer.* 1999, **8**:665-668
- Kawamori T, Tanaka T, Hirose M, Mori H: **Inhibitory effects of d-limonene on the development of colonic aberrant crypt foci induced by azoxymethane in F344 rats.** *Carcinogenesis.* 1996, **17**:369-372
- Gelb MH, Tamanoi F, Yokoyama K, et al: **The inhibition of protein prenyltransferases by oxygenated metabolites of limonene and perillyl alcohol.** *Cancer Letters.* 1995, **91**:169-175
- Elegbede JA, Maltzman TH, Elson CE, et al: **Effects of anticarcinogenic monoterpenes on phase II hepatic metabolizing enzymes.** *Carcinogenesis.* 1993, **14**:1221-1223
- Hardcastle IR, Rowlands MG, Barber AM, Grimshaw RM, Mohan MK, Nutley BP, Jarman M: **Inhibition of protein prenylation by metabolites of limonene.** *Biochemical Pharmacology.* 1999, **57**(7):801-809
- Berkarda B, Koyuncu H, Soybir G, Baykut F: **Inhibitory effect of Hesperidin on tumour initiation and promotion in mouse skin.** *Research in Experimental Medicine.* 1998, **198**:93-99
- Guthrie N, Carroll KK: **Inhibition of mammary cancer by citrus flavonoids.** *Advances in Experimental Medicine & Biology.* 1998, **439**:227-236
- Hakim IA, Harris RB, Ritenbaugh C: **Citrus Peel Use Is Associated With Reduced Risk of Squamous Cell Carcinoma of the Skin.** *Nutrition & Cancer.* 2000, **37**(2):43-50
- Hakim IA, Harris RB, Weisgerber U: **Tea intake and skin squamous cell carcinoma: influence of type of tea beverages.** *Cancer Epidemiology, Biomarkers and Prevention.* 2000, **9**:727-731
- Waksberg J: **Sampling methods for random digital dialing.** *J. Am. Stat. Assoc.* 1978, **56**:33-43
- Nutrition Data System, (NDS) version 2.9;** Nutrition Coordinating Center, University of Minnesota, 1997
- Stata Corp.** Stata statistical software, intercooled stata, release 6.0, College Station, TX: Stata Corporation, 1999
- He L, MO H, Hadisusilo S, Qureshi AA, Elson CE: **Isoprenoids suppress the growth of murine B16 melanomas in vitro and in vivo.** *Journal of Nutrition.* 1997, **127**(5):668-674
- McCarty MF: **Selenium, calcium channel blockers, and cancer risk – the Yin and Yang of apoptosis? [Review]** *Medical Hypotheses.* 1998, **50**(5):423-433
- Uedo N, Tatsuta M, Iishi H, Baba M, Sakai N, Yano H, Otani T: **Inhibition by D-limonene of gastric carcinogenesis induced by N-**

- methyl-N'-nitro-N-nitrosoguanidine in Wistar rats. *Cancer Letters* 1999, **137(2)**:131-136**
36. Conney AH, Lu Y, Lou Y, Xie J, Huang M: **Inhibitory effect of green and black tea on tumor growth.** [Review] *Proceedings of the Society for Experimental Biology & Medicine* 1999, **220(4)**:229-233
 37. Chen NY, Ma WY, Yang CS, Dong Z: **Inhibition of arsenite-induced apoptosis and AP-1 activity by epigallocatechin-3-gallate and theaflavins.** *Journal of Environmental Pathology, Toxicology & Oncology* 2000, **19(3)**:287-295
 38. McNamee D: **Limonene trial in cancer.** *Lancet* 1993, **342**:801
 39. Vigushin DM, Poon GK, Boddy A, English J, Halbert GW, Pagonis C, Jarman M, Coombes RC: **Phase I and pharmacokinetic study of D-limonene in patients with advanced cancer.** *Cancer Chemotherapy. Pharmacol* 1998, **42**:111-117
 40. Franke AA, Cooney RV, Custer LJ, Mordan LJ, Tanaka Y: **Inhibition of neoplastic transformation and bioavailability of dietary flavonoid agents.** *Advances in Experimental Medicine & Biology* 1998, **439**:237-248
 41. Saija A, Tomaino A, Trombetta D, Giacchi M, De Pasquale A, Bonina F: **Influence of different penetration enhancers on in vitro skin permeation and in vivo photoprotective effect of flavonoids,** *International Journal of Pharmaceutics, Volume* 1998, **175(1)**:85-94
 42. Kooststra A: **Protection from UV-B-induced DNA damage by flavonoids.** *Plant Mol. Biol* 1994, **26**:771-774
 43. Tewari S, Gupta V, Bhattacharya S: **Comparative study of antioxidant potential of tea with and without additives.** *Indian Journal of Physiology & Pharmacology* 2000, **44(2)**:215-219

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMedcentral will be the most significant development for disseminating the results of biomedical research in our lifetime."

Paul Nurse, Director-General, Imperial Cancer Research Fund

Publish with **BMC** and your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours - you keep the copyright



Submit your manuscript here:

<http://www.biomedcentral.com/manuscript/>

editorial@biomedcentral.com